

Endometriosis Awareness Month: Enduring “Endo”

March is Endometriosis Awareness Month. According to the American College of Obstetricians and Gynecologists (ACOG), endometriosis is when the tissue that forms the lining of the uterus (the endometrium) is found outside the uterus, causing bleeding, inflammation, and scarring.

While the most common symptom of endometriosis is pelvic pain, this is also a common side effect women experience during their menstrual cycle. In fact, according to the National Institutes of Health (NIH), more than 80% of women experience pelvic pain during their period. This may make it difficult to differentiate what is causing your pain – especially because the pain caused by endometriosis typically occurs during the same time.

Dr. Vanessa Foster, an OSF HealthCare OB/GYN, recommends talking to your OB/GYN if you experience heightened pain during your menstrual cycle, as your provider can help determine its root cause.

Dr. Vanessa Foster, OB/GYN, OSF HealthCare

“Sometimes it is difficult to tell, because everybody’s pain threshold and how they tolerate pain is different. But usually, endometriosis pain is associated with your menstrual cycle. Don’t ignore the pain. As women, we get busy in our lives and are taking care of everything else and you may just ignore it and take Tylenol or Ibuprofen – but I would just say don’t ignore it.” (:24)

She adds that not all women who experience pelvic pain have endometriosis. Sometimes, only a small amount of endometriosis may cause severe, debilitating pain while in other cases, a higher amount of endometriosis may cause little pain. In fact, many women with endometriosis may have no symptoms at all.

In some cases, endometriosis is not detected until a woman is at a stage in her life when she is trying to start a family.

Dr. Vanessa Foster, OB/GYN, OSF HealthCare

“Endometriosis affects 5-10% of reproductive age women, and about a third of them will have infertility problems associated with the endometriosis.” (:12)

Because of this, if you have been struggling with infertility, Dr. Foster recommends talking to your health care provider about the possibility of having endometriosis that has gone undiagnosed.

In addition to being a woman of reproductive age, the NIH adds that other risk factors for endometriosis can include having a mother, sister, or daughter with endometriosis; starting your menstrual period at an early age (before age 11), having short monthly cycles (less than 27 days), or having heavy menstrual periods that last more than seven days.

Dr. Foster says that while most OB/GYN providers can typically determine the presence of endometriosis during an examination, further steps need to be taken to know for sure.

Dr. Vanessa Foster, OB/GYN, OSF HealthCare

“It can only be diagnosed surgically. So you have to look at the endometriosis and take a biopsy of it. A lot of times, we can diagnose it based on history and everything falls into the endometriosis category – but to definitively diagnose it, you need to have surgery.” (:19)

If you think you might have endometriosis, Dr. Foster recommends making an appointment with your OB/GYN and bringing a journal of some sort with you to your appointment to share with your provider.

Dr. Vanessa Foster, OB/GYN, OSF HealthCare

“The best thing is to keep a menstrual calendar. We want to know when the pain started, how long the pain lasts, what it is associated with, does it come before the onset of bleeding or does it come after. So the best advice I have is to keep a very detailed menstrual calendar.” (:21)

To find an OB/GYN provider near you, go to [osfhealthcare.org](https://www.osfhealthcare.org).